

DZHAGATSPANYAN, A.K., otv.red.; ALLAKHVERDYAN, G.O., otv.red.;
SLXUNI, A.G., red.isd-va; KAPLANYAN, M.A., tekhn.red.

[Specialization and cooperation in the machinery industry]
Spetsializatsiya i kooperirovanie v mashinostroenii. Erevan,
1959. 144 p. (MIRA 12:8)

1. Akademiya nauk Armyanskoy SSR, Yerevan. Institut ekonomiki.
(Machinery industry)

DZHAETSPANYAN, R. V.

V1391. Mass-spectrometric method of analysis of chlorine derivatives of methane. S. E. Kuprivanov, R. V. Dzhagetspanyan, M. V. Tikhomirov, and N. H. Tunitskiy (L. Ya. Karpov Physico-Chem. Scientific Research Inst). Zaved. Lab., 1955, 21, No. 10, p. 1182-1188.

Methods for the analysis of mixtures of methane, chloromethane, dichloromethane, chloroform and carbon tetrachloride are described.

G. S. Smith

DZHAGATSPANYAN, R.V.; TUNITSKIY, N.N.

~~CONFIDENTIAL~~

Experimental study of the dilution of chromatographic bands.
Dokl.AN SSSR 105 no.6:1282-1284 D '55. (MLRA 9:4)

1.Fiziko-khimicheskiy institut imeni L.Ya.Karpeva. Predstav-
lene akademikom V.A.Karginym.
(Chromatographic analysis)

DZHAGATSPANYAN, R. V.

PHASE I BOOK EXPLOITATION 300/2386

Россов, Яззико-Кхлалчешекы Инатену

Problemy fizicheskoy khimii, ch. 2, vyp. 2 (Problems in Physical Chemistry: Transactions of the Institute, no. 2), Moscow, Goskhimizdat, 1959. 202 p., 1,000 copies printed.

Editorial Board: Ya. M. Vashchavsky, Doctor of Chemical Sciences; O. S. Zhurav, Doctor of Chemical Sciences; V. A. Kargin, (Moscow); V. M. Kozlovskiy, Doctor of Chemical Sciences (Kiev, U.S.S.R.); V. S. Medvedev, Assistant; S. Ya. Pishchinskii, Doctor of Chemical Sciences; V. M. Chepurchenko, Candidate of Chemical Sciences; V. D. Chavakova (Editorial Secretary); Candidate of Chemical Sciences; Ed.: I. A. Vashukov; tech. Ed.: Ye. O. Shpak.

PURPOSE: This collection of articles is intended for physical chemists.

CONTENTS: The collection is the second issue of the Transactions of the Scientific Research Institute of Physical Chemistry named L. Ya. Karpov. It contains 17 articles which review

Tenklin, M. I., M. M. Morozov, V. M. Piznev (Deceased), M. O. Apol'shchik, V. I. Lukyanova, and Y. A. Dvorkin. The Oxidation of Anomols Over a Nonplatinum Catalyst. 144

2. Yamashita, S. T., S. A. Zamaebakura, T. S. I. Orliva, A. V. Gerasimov, N. K. Kuroyan, L. N. Pospelov, A. A. Igin, Y. K. Il'inskiy, N. A. Silyavskaya, and V. I. Chudakovskiy. Kinetics of decomposition, and the explosion of carbon tetrachloride, CCl_4 (Japan). How to find the kinetic equation of a reversible reaction. 39

Kolot'yin, Ya. M. The Effect of the Specific Adsorption of Ions on the Kinetics of Hydrogen Evolution and the Structure of the Metal-Solution Boundary

Verkhavsky, Ya. M. The Nature and Mechanism of Electro-
philic Nitrogen Exchange

Krishna, Z. V. Crystallochemical Data on the Nature of
the Mutual Effect of Atoms 97

WATKINS, D. J., Investigation of the Effect of Intramolecular Interaction on the Ultraviolet Absorption Spectra of Aromatic Compounds 107

118
S. I. V. S. Kuev and B. P. Ormont. Investi-
gation of Hydrolysis in the System Nitrobenzene at
High Temperatures and the Dependence of the Free Energy
of C-Ng Formation on Its Composition and Structure

Shcherbak, A. G.; M. A. Dobronogovskiy, A. A. Dostoyev, L. I. Zaitseva and M. S. Rykova. Study of the Field of Forces of Damages from a Cylindrical Impactor with CO₂ as a Power-Cell Source of Y Ablation. *1989*

BOGOMOL'Y, V. K.; B. G. VASIL'YEV and N. M. GUNTAKSKY. Study of the Ionization and Dissociation of n-Octane and n-Nonane Radical Ions by the Method of Bombardment with "Quasi-Resonant" Electrons

ESKEND, A. E. Radiation-Chemical Effects in Solid
Inorganic Salts 161

Polozhenko, V. A., Ye. V. Burdakov, and L. I. Yatskovskaya.
Kinetics of the Process of Benzene Oxidation in an Aqueous
Solution of Potassium Permanganate. *Dokl. Akad. Nauk SSSR*,
1977, 245, No. 1, 169-171. *Chem. Abstr.*, 1977, 86, 129163d.
Russian-O English Translation of the Original.

197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

University, T. A. and D. A. Golden. The Problem of the Phase Composition of the System $\text{H}_2\text{O}-\text{NaOH}-\text{KOH}$ at Low Temperatures

Gerasimov V. D. and A. A. Zarnokhova. Constitution of the
Enzyme-Induced Oxidation of Leuciform Dyes

5(3)

SGV/64-59-4-7/27

AUTHORS: Urin, M. G., Dzbagatspanyan, R. V.

TITLE: Quantitative Analysis of the Commercial Product DDT by Means of Infrared Spectrometry (Kolichestvennyy analiz tekhnicheskogo produkta DDT s pomoshch'yu infrakrasnoy spektrometrii)

PERIODICAL: Khimicheskaya promyshlennost', 1959, Nr 4, pp 27 - 28 (USSR)

ABSTRACT: Among the 15 compounds which were found in the commercial DDT the n,n'-isomer DDT (I) has the strongest toxic effect. The spectroscopic method allows an especially precise, simple, and rapid quantitative determination of this isomer. The commercial DDT may be regarded as a mixture of the (I) and o,n'-isomer DDT (II) in this connection the n,n'-isomers DDD and DDKh are also determined in the former. A spectroscopic method was elaborated for the quantitative analysis of two-, three- and four-component-mixtures of the last mentioned substances. The experiments were carried through by means of the spectrometer IKS-11 with a Nernst needle as radiation source. Carbon disulphide was used as solvent. The following analytical wave lengths were used: for (I) 12.3 μ , for (II) 9.6 μ , for the n,n'-isomer DDD 12.4 μ and for the n,n'-isomer DDKh 10.2 μ . 16

Card 1/2

Quantitative Analysis of the Commercial Product DDT by SOV/64-59-4-7/27
Means of Infrared Spectrometry

absorption coefficients K_{ij} (Table 1) were determined and two-, three- and four-component standard mixtures of a similar composition as the commercial DDT-product were analysed (Table 2). The measuring accuracy for (I) and (II) is given with a mean error of 1 and 2.5%. There are 2 figures, 2 tables, and 15 references, 4 of which are Soviet.

Card 2/2

21.3000

77278
SOV/63-4-6-12/37

AUTHORS: Dzhagatspanyan, R. V., Zetkin, V. I., (Candidates of Chemical Sciences)

TITLE: Concerning Some Radiochemical Reactions of Organic Synthesis

PERIODICAL: Khimicheskaya nauka i promyshlennost', 1959, Vol 4, Nr 6, pp 761-769 (USSR)

ABSTRACT: This is a review of literature data concerning the use of nuclear radiation for the initiation of chain and nonchain reactions; and utilization of reactions initiated by nuclear radiation in the chemical industry. The use of Co^{60} and Cs^{134} (as well as wastes of nuclear reactors) as the sources of γ -radiation initiating many chemical reactions was discussed. The following processes were considered in detail: chlorination and bromination of organic compounds initiated by radiation. Chlorination of benzene initiated by Co^{60} in the absence of O_2 with "Freon-12" (CF_2Cl_2) as a diluent

Card 1/5

Concerning Some Radiochemical Reactions
of Organic Synthesis

77278
SOV/63-4-6-12/37

lead to the conclusion that there is a critical temperature for each benzene: chlorine: freon ratio in the reacting mixture. At the freezing point the yield of hexachlorocyclohexane and its isomers sharply decreases (to 0.75-1.5 mole/100 ev). A further lowering of the temperature results in a sharp increase in the yield of hexachlorocyclohexane (1,500 mole/100 ev). Thus it was suggested that a chain reaction takes place on the surface and in the crystals of benzene. Chlorination and bromination in the presence of radioactive initiators is conducted at a lower temperature and is safer than catalytic and chlorination under ultraviolet irradiation. Addition of silanes to unsaturated compounds, initiated by γ -radiation, sulfochlorination, and sulfoxidation are discussed. Chlorination of trimethylchlorosilane and ethyltrichlorosilane with elemental chlorine, initiated by

Card 2/5


Concerning Some Radiochemical Reactions
of Organic Synthesis

77278
SOV/63-4-6-12/37

γ -radiation, proceeds 100 times faster than chlorination with ultraviolet irradiation. The yield is 90%. The scheme shown in Fig. 2 of industrial chlorination of alkylchlorosilanes initiated by nuclear radiation is suggested. There are 84 references, 39 U.S., 6 U.K., 2 Swiss, 5 German, 3 French, 29 Soviet. The 5 most recent U.S. and U.K. references are: P. Cheek, V. S. Leinnenbom, J. Phys. Chem., 62, Nr 72, 1475 (1958); R. A. Cox, A. J., Swallow, J. Chem. Soc., 3727; (1958); J. F. Black, E. F. Baxter, Soap and Chem. Specialties, 34, Nr 10, 43 (1958); A. M. El. Abbady, L. C. Anderson, J. Am. Chem. Soc., 80, 1737 (1958); E. T. McBee, C. W. Roberts, G. W. R. Puerckhauer, J. Am. Chem. Soc., 79, 484 (1957).

Card 3/5

77278, SOV/63-4-6-12/37



Card 4/5

Concerning Some Radiochemical Reactions
of Organic Synthesis

77278
SOV/63-4-6-12/37

Fig. 2. Flow sheet of radiochemical chlorination of alkylchlorosilanes: (1) raw materials storage tank; (2) mixer; (3) heat exchangers; (4) reactor; (5) well for keeping the initiator (source of γ -radiation); (6) shielding; (7) HCl evaporator; (8) steam coil; (9) drop trap; (10) heaters; (11) fractionating columns; (12) storage for the unreacted alkylchlorosilane; (13) storage for the manufactured product; (14) perchlorinated products; (15) raw material pump; (16) HCl exit.

Figure on Card 4/5

Card 5/5

S/882/62/000/002/041/100

A057/A126

AUTHORS: Dzhagatspanyan, R.V., Zetkin, V.I., Zykova, Ye.N., Filippov, M.T.

TITLE: A method for the sulfochlorination of polyethylene

SOURCE: Sbornik izobreteniy; plastmassy i sinteticheskiye smoly. no. 2.
Kom. po delam izobr. i otkrytiy. Moscow, TsBTI, 1962, 25 [Author's
certificate no. 128142, cl. 39c, 2501 (appl. no. 637114 of August
24, 1959)]

TEXT: To accelerate sulfochlorination and to avoid a simultaneous destruc-
tion of polyethylene the following procedure is suggested: the 3% polyethylene
solution diluted in carbon tetrachloride is treated with a mixture of gaseous
sulfur dioxide and chlorine (2 : 1) at 70° C and is Gamma irradiated by a Co⁶⁰
source with an activity of 1,400 g.equiv. radium, and a dosis capacity of 120 r/
/sec for 15 - 40 sec. SO₂ and Cl₂ are introduced with a rate of 256 and 178 ml/
/min, respectively. The obtained polymer contains one SO₂Cl group per approxi-
mately 38 methylene groups.

[Abstracter's note: Complete translation]

Card 1/1

5(3)

AUTHORS:

Bardenshteyn, S. B.,
Dzhagatspanyan, R. V., Zetkin, V. I.

S/032/60/026/02/018/057
B010/B009

TITLE:

Analysis of a Mixture of Isomeric Trichlorobenzenes¹ and of a
Mixture of Isomeric Tetrachlorobenzenes by Means of Infrared
Absorption Spectra

PERIODICAL:

Zavodskaya laboratoriya, 1960, Vol 26, Nr 2, pp 167 - 171 (USSR)

ABSTRACT:

A determination of the composition of a six component mixture consisting of tri- and tetrachlorobenzenes cannot be carried out spectroscopically due to the insufficient resolving power of the IKS-11 spectrometer. For this reason it is recommended to separate the mixture into fractions of the isomers of the tri- and tetrachlorobenzenes prior to analysis. In the present paper the 6.97μ absorption band was used as the analytical line for 1,2,3-trichlorobenzene. Carefully cleaned preparations were used (Table 1, Figs 1,2) for determining the spectra of the isomers. Carbon tetrachloride was used as solvent. The sample was separated into the isomers by vacuum distillation. It was found experimentally that the Lambert-Beer law holds for 1,3,5- and 1,2,4-trichlorobenzenes as well as for all tetra-

Card 1/2

Analysis of a Mixture of Isomeric Trichlorobenzenes S/032/60/026/02/018/057
and of a Mixture of Isomeric Tetrachlorobenzenes by B010/B009
Means of Infrared Absorption Spectra

chlorobenzene isomers up to optical densities of 0.4-0.5, which corresponds to the necessary range of concentrations. 1,2,3-trichlorobenzene has to be determined graphically in the case of concentrations of more than 12% by weight. Data concerning the analytical results as well as the repeatability and calibration solutions are given (Table 2). Two samples of commercial products were analyzed by the above method, and the results were compared with those obtained by vacuum distillation. (Table 3). There are 2 figures, 3 tables, and 5 references, 1 of which is Soviet.

Card 2/2

KOLBASOV, V.I., BARDENSHTEYN, S.B., DZHAGATSPANYAN, R.V.

Quantitative analysis of a mixture of monochlorides, based
on their infrared absorption spectra. Zav.lab. 26 no.5:587-
590 '60. (MIRA 13:7)

(Chlorides--Spectra)

FLISSKIY, M.M.; VESELOVSKAYA, I.Ye.; IZHAGATSPANYAN, R.V.

Destruction of graphite anodes in the electrolysis of sodium chloride
in the presence of sulfate ions. Zhur. prikl. khim. 33 no.8:1901-1903
Ag '60. (MIRA 13:9)

(Electrodes, Carbon)

(Electrolysis)

(Salt)

KOLBASOV, V.I.; BARDEISHTEYN, S.B.; DZHAGATSPANYAN, R.V.

Quantitative determination of impurities in industrial epichlorohydrin
from infrared absorption spectra. Zav.lab 26 no.10:1120-1122 '60.
(MIRA 13:10)

(Epichlorohydrin--Spectra)

DZHAGATSPANYAN, R V

137

PHASE I BOOK EXPLOITATION

SOV/5486

Vsesoyuznoye soveshchaniye po vnedreniyu radioaktivnykh izotopov i yadernykh izlucheniya v narodnoye khozyaystvo SSSR. Riga, 1960.

Radioaktivnyye izotopy i yadernyye izlucheniya v narodnom khozyaystve SSSR; trudy soveshchaniya v 4 tomakh. t. 1: Obshchiye voprosy primeneniya izotopov, pribory s istochnikami radioaktivnykh izlucheniya, radiatsionnaya khimiya, khimicheskaya i neftepererabatyvayushchaya promyshlennost' (Radioactive Isotopes and Nuclear Radiations in the National Economy of the USSR; Transactions of the Symposium in 4 Volumes. v. 1: General Problems in the Utilization of Isotopes; Instruments With Sources of Radioactive Radiation; Radiation Chemistry; the Chemical and Petroleum-Refining Industry) Moscow, Gosoptekhizdat, 1961. 340 p. 4,140 copies printed.

Sponsoring Agency: Gosudarstvennyy nauchno-tekhnicheskiy komitet Soveta Ministrov SSSR, and Gosudarstvennyy komitet Soveta Ministrov SSSR po ispol'zovaniyu atomnoy energii.

Ed. (Title page): N.A. Petrov, L.I. Petrenko and P.S. Savitskiy; Eds. of this Vol.: L.I. Petrenko, P.S. Savitskiy, V.I. Sinitsin, Ya. M. Kolotyarkin, N.P. Syrma and R.F. Romm; Executive Eds.: Ye. S. Levina and B. F. Titskaya; Tech. Ed.: E.A. Mukhina.

Card 1/12

137

Radioactive Isotopes (Cont.)

80V/5486

PURPOSE: The book is intended for technical personnel concerned with problems of application of radioactive isotopes and nuclear radiation in all branches of the Soviet economy.

COVERAGE: An All-Union Conference on problems in the introduction of radioactive isotopes and nuclear radiation into the national economy of the Soviet Union took place in Riga on 12-16 April 1960. The Conference was sponsored by: the Gosudarstvennyy nauchno-tekhnicheskii komitet Soveta Ministrov SSSR (State Scientific and Technical Committee of the Council of Ministers, USSR); Glavnoye upravleniye po ispol'zovaniyu atomnoy energii pri Sovete Ministrov SSSR (Main Administration for the Utilization of Atomic Energy of the Council of Ministers, USSR); Academy of Sciences, USSR; Gosplan USSR; Gosudarstvennyy komitet Soveta Ministrov SSSR po avtomatizatsii i mashinostroyeniyu (State Committee of the Council of Ministers, USSR, for Automation and Machine Building) and the Council of Ministers of the Latvian SSR. The transactions of this Conference are published in four volumes. Volume I contains articles on the following subjects: the general problems of the Conference topics; the state and prospects of development of radiation chemistry; and results and prospects of applying radioactive isotopes and nuclear radiation in the petroleum refining and chemical industries. Problems of designing and manufacturing instruments which contain sources of radioactive radiation and are used for checking and automation of technological processes are examined, along with problems of accident prevention in their use. No personalities are mentioned. References accompany some of the articles.

Card 2/12

Radioactive Isotopes

SOV/5486

- Dzhagatspanyan, R.V., V.I. Zetkin, Ye. N. Zyкова, and M.T. Filippov.
Sulfochlorination and Sulfooxidation of Polyethylene, Polypropylene,
Polyisobutylene, and Paraffin Hydrocarbons by Radiation 201
- Topchiyev, A.V., L.S. Polak, N. Ya. Chernyak, V. Ye. Glushnev,
P. Ya. Glazunov, I.V. Vereshchinskiy, N.P. Syrkus, A. Kh. Breger,
and B.I. Vaynshteyn. Radiation-Thermal Hydrocarbon Cracking 206
- Beer, A.A., M.A. Besprozvanny, T.I. Limanova, and M.T. Filippov.
Radiation-Chemical Telomerization 211
- Zimakov, P.V., Ye. V. Volkova, A.V. Fokin, A.D. Sorokin, and V.M.
Belikov. Utilization of Nuclear Radiation Energy in Fluorinated
Olefin Polymerization Processes 219
- Breger, A. Kh., V.B. Osipov, and V.A. Gol'din. Universal Installation
(K-60000) With Co⁶⁰ γ-Radiation Source of 60,000 Gram-Equivalent
Radium Activity, for Simulating Radiation-Chemical Apparatus and
Carrying Out Investigations 227

Card:8/12

S/019/61/000/003/053/101
A154/A027

AUTHORS: Dzhagatspanyan, R.V., Zetkin, V.I., and Zykova, Ye.N.

TITLE: A Method of Sulfo-oxidizing Polyethylene

PERIODICAL: Byulleten' izobreteniy, 1961, No. 3, p. 49

TEXT: Class 39c, 25₀₁. No. 135639 (667154/23 of May 18, 1960).
A method of sulfo-oxidizing (sul'fookisleniya) polyethylene by initiated
reaction of a solution or suspension of polyethylene in an inert solvent with
sulfurous anhydride and oxygen, distinguished by the fact that the process
is initiated by γ -irradiation with the use of Co⁶⁰ as a radioactive source.

✓

Card 1/1

S/019/60/000/009/020/071
A152/A029

AUTHORS: Dzhagatspanyan, R.V.; Zetkin, V.I.; Zykova, Ye.N.; Filippov, M.T.

TITLE: A Method of Sulfochlorinating Polyethylene 1

PERIODICAL: Byulleten' izobreteniy, 1960, No. 9, p. 39

TEXT: Class 39c, 2501. No. 128142 (637114/23 of August 24, 1959). A method of sulfochlorinating polyethylene with sulfur dioxide gas and chlorine. It has the following special feature: the reaction is accelerated by means of conducting the process under the influence of Co⁶⁰ gamma-radiation.

Card 1/1

Dzhagatspanyan, R.V.

~~25(1)~~ 5.3700(B)

00354

S/019/60/000/02/038/221
D031/D005

AUTHORS: Dzhagatspanyan, R.V., Filippov, M.T., Motsarev, G.V.,
Zetkin, V.I., and Koreshkova, N.G.

TITLE: A Method of Obtaining Alkyl and Aryl Polysiloxanes ¹ Containing Chlorine

PERIODICAL: Byulleten' izobreteniy, 1960, Nr 2, p 14 (USSR)

ABSTRACT: Class 12o, 26⁰³. Nr 125564 (624019/23 of 2 April 1959).
This method is for obtaining alkyl and aryl polysiloxanes
containing chlorine by chlorination of the alkyl or aryl
polysiloxanes with elementary chlorine. To increase the out-
put, the process is carried out when the basic material is
initiated by γ radiation of radioactive cobalt at a tempera-
ture of about 0°C. X

Card 1/1

5(1)

S/019/59/000/23/046/432
D046/D007

AUTHORS: Dzhagatspanyan, R.V., Kokhanov, G.N., Veselovskaya, I.Ye. and
Mulin, Ye.V.

TITLE: A Metal Current Lead to the Carbon ⁶Graphite ⁷Anodes for Electrolyzers

PERIODICAL: Byulleten' izobreteniy, 1959, Nr 23, p 14 (USSR)

ABSTRACT: Class 12h, 2. Nr 124423 (611988/23 of 18 November 1958). With addition of claim Nr 617444. To reduce voltage resistance loss in the anodes, the lead is bimetallic. The outer sheeting is made by metal producing a closing layer at the anode polarization.

Card 1/1

5(1)

SOV/19-59-6-29/309

AUTHOR: ~~Dzhagatspanyan~~, R.V., Zetkin, V.I., and Maksimov, M.P.

TITLE: A Method of Chlorinating Alkyl Chlorosilanes with Chlorine

PERIODICAL: Byulleten' izobreteniy, 1959, Nr 6, p 10 (USSR)

ABSTRACT: Class 12o, 26₀₁. Nr 118500 (581222 of 30 July 1957).
To increase the output and the speed of reaction, and to decrease the temperature, the reaction zone is being irradiated with γ -rays.

Card 1/1

54600

53700

2209

24372
S/063/61/006/002/004/004
A105/A129

AUTHORS: Dzhagatspanyan, R. V., Zetkin, V. I., Filippov, M. T.

TITLE: A study of the radiation addition of silicochloroform to certain olefines

PERIODICAL: Zhurnal vsesoyuznogo khimicheskogo obshchestva im. D. I. Mendeleyeva, v. 6, no. 2, 1961, 231 - 232

TEXT: The authors refer to the reaction of addition of hydridesolanes at the multiple bonds of organic compounds as being the most promising process for the production of various silicon-organic monomers of high purity. The present article deals with an investigation of the addition of trichlorosilane to nonene-1 and cyclohexene under the action of γ - radiation at two dose rates (70 r/sec and 123 r/sec), 25 and 70°C and a molar ratio of the reagents of 1:1. Co^{60} was the radiation source having an activity of 1,400 g-equiv. of radium. The reaction products were decomposed by fractional distillation. The formed compounds were identified by the chlorine content and the determination of carbon and hydrogen could be carried out by transforming the halidesilanes into the corresponding alkoxy-derivatives. Tables 1 and 2 show the results of the in-

Card 1/5

24372

S/063/61/006/002/004/004

A study of the radiation addition of silicochloroform.. A105/A129

vestigations of the addition of trichlorosilane to nonene-1. It is seen that an increase in the dosage from 70 r/sec to 123 r/sec increases the rate of reaction. Table 3 lists the results of the addition of trichlorosilane to cyclohexene. The values of G calculated from the experimental data obtained by the present authors in the reaction of addition of trichlorosilane to nonene-1 and cyclohexene were found to be comparatively low and close to the values of other experiments (Ref. 4). Under the effect of γ -emission the addition of trichlorosilane to nonene was found to take place with about the same yield as when initiated with benzoyl peroxide viz. 60.9% at 85°C in the presence of weight 3% of benzoyl peroxide (Ref. 5). There are 3 tables and 5 references: 2 Soviet-bloc and 3 non-Soviet-bloc.

Table 1: 1 - Addition of trichlorosilane to nonene-1. Dose rate 70 r/sec, temperature 25°C, mol. ratio 1:1; 2 - reagent charge; 3 - radiation time h; 4 - yield, % of theoretical; 5 - radiation-chemical yield G (mol/sec 100 ev); 6 - nonene-1; 7 - SiHCl_3 .

Card 2/5

9.4160 (also 2801)

20704
S/120/61/000/001/042/062
E192/E382

11.8100
AUTHORS: Dzhagatspanyan, R.V., Maksimov, M.P. (Deceased)
and Yesel'son, M.P.

TITLE: A Device for Recording High-speed Processes

PERIODICAL: Pribory i tekhnika eksperimenta, 1961, No. 1,
pp. 132 - 137

TEXT: Continuous recording of the changes of infra-red spectra during chemical reactions is of considerable practical importance but very often the speed of response of the infra-red receivers (wavelengths from 2 - 20 μ or more) is not fast enough to give a suitable resolving time. A faster device was proposed by Bonch-Bruyevich and Imas (Ref. 1) in 1955 but it resulted in the deterioration of the signal-noise ratio of the receiver. In the following a high-speed recording instrument with an amplifier furnished with a bolometer inertia-correcting circuit is described. The recording speed for a single spectrum can be as high as 10^{-2} or 10^{-3} sec, the resolving time being 10^{-4} to 10^{-5} sec. The instrument can work as a pyrometer (recording rapid changes of the incident

Card 1/5

20704

S/120/61/000/001/042/062
E192/E382

A Device for Recording

thermal radiation) or it can perform measurements at a chosen point of the spectral range. When operating as a spectrometer the instrument can register infra-red spectra $I = f(\lambda)$ of various materials at speeds from $1 - 10^{-5}$ sec. The limiting resolving time of the instrument is 6×10^{-3} which is equivalent to 10^{-4} sec. Consequently, by recording a successive range of spectra it is possible to plot the spectrum $I = \varphi(\lambda, t)$ of a chemical reaction. When used as a pyrometer the instrument can register rapid changes of the radiation intensity as a function of time ($I = f(t)$) at speeds ranging from $1 - 10^{-3}$ sec, the limiting resolving time being of the order of 10^{-4} sec. This is necessary in solving various engineering problems such as the investigation of explosions, rapid combustion, etc. In both types of operation a calibrating voltage curve is recorded simultaneously with the curves representing the investigated processes; the calibration curve permits determination of the duration of the

Card 2/5

20704

S/120/61/000/001/042/062
E192/E382

A Device for Recording

process and its significant sections. The spectra are recorded by means of a long-persistence cathode-ray tube (oscillograph ENO-1 (ENO-1)). The instrument consists of the following units: 1) a pre-amplifier which is situated in the immediate vicinity of the receiver; 2) the main amplifier, provided with an inertia-correction circuit; 3) an electronic switch having two inputs and operating at the frequency of 200 kc/s; 4) a calibration oscillator producing sinusoidal waveforms having frequencies of 10, 50, 100, 500 and 1 000 c.p.s; 5) a control pulse generator giving rectangular pulses repeated at a frequency of 50 c.p.s; 6) a synchronisation amplifier, permitting synchronisation of the time base by a positive or negative internal signal; 7) a triggered time base operating at frequencies 1, 3, 10, 30, 100, 300 and 1 000 c.p.s. and power supplies comprising high-voltage and low-voltage rectifiers and an electronic stabiliser having a stabilisation coefficient of 1200. The main amplifier consists of an input cathode follower, a two-stage amplifier furnished with high- and low-frequency correction circuits and an inertia-correcting circuit which is separated

Card 3/5

20704

S/120/61/000/001/042/062
E192/E382

A Device for Recording

from the amplifier by means of a cathode follower. The bandwidth of the amplifier at 3 db extends from 1 c.p.s. to 0.8 Mc/s. The inertia-correction is achieved by means of an RC network. The range of the time constants of the receivers is divided into the following sub-ranges: 1-3, 3-8, 5-15, 10-30 and 30-100 μ s. The desired sub-range is provided by switching-in a suitable capacitance, while the continuous control is achieved by varying the resistance. The electronic switch consists of a multivibrator operating at a frequency of 200 kc/s, a limiter and two controlled tubes. The switch is perhaps unusual in that a cathode-coupled feedback stage is used as the limiter. The calibration oscillator is based on the usual RC circuit. The synchronisation amplifier consists of two tubes, while the time-base generator comprises two multivibrators, a charging pentode and a cathode follower. The time constant of the correction network should be made equal to the time constant of the receiver before the instrument can be used in measurements. This is done by introducing a

Card 4/5

20704

S/120/61/000/001/042/062
E192/E382

A Device for Recording

rotating (modulating) disc between the source and the receiver, the purpose of the disc being to produce light pulses having a sharp leading edge. The instrument was tested as a pyrometer, a photoresistor having a time constant of 40 μ s being used as a receiver. There are 4 figures and 1 Soviet reference.

SUBMITTED: December 28, 1959

X

Card 5/5

21135

S/19C/61/003/004/01C/C14
B101/B207

15 2116

22-02-77

AUTHORS: Dzhagatsoyan, R. V., Zetkin, V. I., Motsarev, G. V.,
Filippov, M. T.

TITLE: Chlorination of organo-silicon monomers and polymers under
the action of gamma rays. I. Chlorination of liquid poly-
phenyl-methyl siloxane and of polydimethyl siloxane rubber.
The infrared spectra of the chlorination products

PERIODICAL: Vysokomolekulyarnyye soyedineniya, v. 3, no. 4, 1961, 607-612

TEXT: In the introduction, the authors state that initiating the chlorina-
tion of organosilicon compounds by means of ultraviolet light proceeds too
slowly, however, that chemical initiators as e.g., benzoyl peroxide re-
quire a higher temperature at which a sufficient chlorination of methyl
chloro silanes is not possible owing to their instability. Therefore, the
present study aimed at initiating chlorination by means of gamma rays of
Co⁶⁰ at low temperatures. The following compounds were chlorinated:

1) Polyphenyl-methyl siloxane (poly-PMS) (molecular weight 2000), and 2)
three samples of polymethyl siloxane rubber (poly-MSR) (molecular weight

Card 1, 5

21135

S/190/61/003/004/010/014
B101/B207

Chlorination of ..

400,000-500,000). 4-5% solutions of the polymers in CCl_4 were used, to which chlorine taken from the cylinder was added. The samples were irradiated in sealed ampoules at 0°C with gamma rays of Co^{60} , activity 1400 G. equivalent. Tables 1 and 2 list the results. Heating with 40% KOH of a chlorinated poly-PMS sample with 55.5% Cl yielded a paste from which crystals with a chlorine content of 64.3-66.7% were separated. On the basis of analytical results, they obtain the empirical formula $\text{C}_6\text{H}_6\text{Cl}_4$ or $\text{C}_6\text{H}_4\text{Cl}_4$.

The infrared spectra of the oily residue of hydrolysis showed an intensive band at $9-10\ \mu$ which corresponds to the Si-O bond. Chlorination of poly-MSR led, according to the sample used, to quite different results with respect to the intensity of reaction and the chlorine content of the product obtained. This is due to impurities (catalyst residues) in commercial poly-MSR. Study of the infrared spectra yielded 3690 and $3615\ \text{cm}^{-1}$ bands both in initial and chlorinated rubber. These bands are due to OH groups ($3690\ \text{cm}^{-1}$ free OH; $3615\ \text{cm}^{-1}$ OH with H bond). Accordingly, commercial poly-MSR contains silanol groups. As a result of spectral analysis the following is stated: though the IR spectra of chlorinated poly-PMS and poly-MSR differ from those of the initial samples, no absorption bands were found to exist which are characteristic of chlorinated substances.

and 2.5

21135

S/190/61/003/004/010/014
B101/B207

Chlorination of ...

There are 2 figures, 5 tables, and 14 references: 8 Soviet-bloc and 6 non-Soviet-bloc. The 2 references to English-language publications read as follows: Ch. Tamborcki, H. W. Post, J. Org. Chem., 17, 1400, 1952; C. W. Joung, P. C. Servais, C. C. Currie, M. J. Hunter, J. Amer. Chem. Soc., 70, 3750, 1948.

SUBMITTED: July 15, 1960

① Опыт №	② Загружено реагентов, г		③ Соотно- шение ① молей Cl основомоль	④ Мощность дозы, р/сек	⑤ Премия облуче- ния, мин	⑥ Вес продукта, г	⑦ Содержание хлора, %	
	② хлор	② полимер					⑦ анал. тело	⑦ анал. тело
1	4,70	4,65	1,97:1	70	30	8,874	48,6	51
2	7,2	4,65	2,98:1	70	30	11,425	59,5	61
3	4,9	3,12	3:1	120	2	8,4819	50,1	61
4	4,9	3,12	3:1	120	5	7,0128	54,0	61
5	4,0	3,12	3:1	120	10	7,8840	58,7	61
6	2,43	4,68	1:1	120	15	7,1914	33,9	34,3
7	3,3	3,12	2,03:1	120	15	5,9615	50,2	51

Card 3/5

2135

S/190/61/003/004/010/014
B101/B207

Chlorination of ...

Table 1. Chlorination of a 4% solution of $[(C_6H_5)(CH_3)SiO]_n$ in CCl_4 at $0^\circ C$.
Legend: 1) No. of the experiment; weighed portion, g; a) chlorine; b) polymer; 3) ratio mole Cl: mole structure unit of polymer; 4) dose, r/sec; 5) time of irradiation; min; 6) weight of the product obtained; 7) chlorine content; c) found; d) calculated on the assumption of chlorine addition to the double bonds of the ring.

1 Эксп. №	2 Реагенты		3 Обра- зец	4 Соотноше- ние, моль Cl основомоль	5 Темпера- тура, °C	6 Мощность дозы, р/сек	7 Время облуче- ния, мин.	8 Содержание хлора в продукте, %	
	а) хлор	б) полимер						в) найде- но	г) рас- считано
1	3,99	3,83	1	1,09:1	40	70	30	32,3	32,8
2	7,77	3,83	1	2,05:1	0	70	15	41,1	50,0
3	7,93	3,83	1	2,17:1	0	70	30	48,9	52,1
4	10,97	3,83	1	3:1	22,5	70	30	49,8	59,2
5	10,97	3,83	1	3:1	0	70	60	55,3	59,2
6	13,2	3,14	2	4,4:1	0	120	10	64,6	67,8
7	2,38	2,49	3	1:1	20	—	—	37,7	32,4
8	4,08	2,49	3	1,64:1	20	—	—	46,6	45,0
9	4,44	1,87	3	2,5:1	0	120	5	60,7	54,2
10	5,28	1,87	3	2,97:1	0	120	5	63,0	58,5

Card 4/5

21135

S/190/61/003/004/010/014
B101/B207

Chlorination of ...

Table 2. Chlorination of a 4% solution of $[(CH_3)_2SiO]_n$ in CCl_4 .

Legend: 1) No. of experiment; 2) weighed portion, g; a) chlorine; b) polymer; 3) No. of sample; 4) ratio mole Cl : mole structure unit of polymer; 5) temperature; 6) dose, r/sec; 7) irradiation time, min; 8) chlorine content in the product; c) found; d) calculated; *) 10 hr in scattered light.

X

Card 5/5

DZHAGATSPANIAN, R.Y.; ZETKIN, V.I.; FIKIPPOV, M.T.

Study of the radiation induced addition of silicochloroform to
some olefins. Zhur. VKHO 6 no.2:231-232 '61. (MIRA 14:3)
(Silane) (Olefins)

S/032/61/027/003/010/025
B101/B203

AUTHORS: Kolbasov, V. I., Bardenshteyn, S. B., and Dzhagatspanyan, R. V.

TITLE: Quantitative analysis of crude trichloro ethane by means of
infrared absorption spectra

PERIODICAL: Zavodskaya laboratoriya, v. 27, no. 3, 1961, 295-296

TEXT: To elaborate an efficient method for the simultaneous production of perchloro-vinyl resin and trichloro ethane it was necessary to analyze the crude trichloro ethane which consisted of 50-60% 1, 2-dichloro ethane, 40-50% 1, 1, 2-trichloro ethane, and 3-5% tetrachloro ethanes. An analysis of the mixture by rectification takes much time (2-3 days) and is independent. The present paper describes a method for the quantitative analysis of crude trichloro ethane on the basis of infrared spectra taken with an MKC-14 (IKS-14) split-beam spectrophotometer. Such an analysis takes only about one hr. The infrared spectra of the substances concerned are described in publications: 1, 2-dichloro ethane (Ref. 1: A. Berton, Chim. analyt. 38, No. 6, 207 (1956); Ref. 2: G. Pirlet, Bull. Soc. chim. belges, 58, No. 1, 28 (1949); Ref. 3: J. K. Brown, H. Sheppard, Trans. Faraday Soc,

Card 1/5

S/032/61/027/003/010/025
B101/B203

Quantitative analysis ...

48, 128 (1952)); 1, 1, 2-trichloro ethane (Ref. 2); 1, 1, 1, 2-tetrachloro ethane (Ref. 2 and Ref. 4: I. R. Nielsen, C. Liang, Z. W. Daasch, J. Opt. Soc. Amer., 43, 1071 (1953)); 1, 1, 2, 2-tetrachloro ethane (Refs. 1, 2, 4), as well as the method for the quantitative determination of their mixtures (Ref. 2, Ref. 5: A. I. Finkel'shteyn, Ts. N. Roginskaya et al., Zavodskaya laboratoriya, XXV, 8, 932 (1959)). The proposed analysis of the quaternary mixture of 1, 2-dichloro ethane, 1, 1, 2-trichloro ethane, 1, 1, 1, 2- and 1, 1, 2, 2-tetrachloro ethane is distinguished from the analysis described in Refs. 2, 5 by the use of the MKC-14 (IKS-14) split-beam spectrometer, and the calculation of concentration on the basis of standard mixtures using the method of least squares (pentachloro ethane and 1, 1, 2, 2-tetrachloro ethane give superimposed bands at 1017 cm^{-1} , and are determined summationally). CCl_4 was used as a solvent. Well purified preparations made by E. Sonin, the constants of which agreed with published data, were employed (Table 1). The figure shows the infrared spectra of the four substances studied (thickness of the absorption layer 0.01 mm). The optical density was determined according to Ref. 6 (Z. Williams, Anal. Chem. 29, No. 10, 1551 (1957)), the concentration of components was calcu-

Card 2/5


S/032/61/027/003/C10/025
B101/B203

Quantitative analysis ...

lated by means of a system of four linear equations. To determine the coefficients of the equations, standard mixtures were prepared for the concentrations required, and the optical density of the analytical absorption bands of the components was determined. A total of 95 binary, ternary, and quaternary mixtures were prepared. The measured results were evaluated by the method of least squares. There are 1 figure, 2 tables, and 9 references: 4 Soviet-bloc and 5 non-Soviet-bloc. [Abstracter's note: Complete translation.]

Legend to the figure:

- 1) 1, 2-dichloro ethane;
- 2) 1, 1, 2-trichloro ethane;
- 3) 1, 1, 2, 2-tetrachloro ethane;
- 4) 1, 1, 1, 2-tetrachloro ethane



Card 3/5

Quantitative analysis ...

S/032/61/027/003/C10/025
B101/B203

Legend to Table 1: 1) Compound, 2) boiling point, 3) specific gravity,
4) refractive index, 5) 1, 2-dichloro ethane, 6) 1, 1, 2-trichloro ethane,
7) 1, 1, 1, 2-tetrachloro ethane, 8) 1, 1, 2, 2-tetrachloro ethane

Legend to Table 2: 1) Compound, 2) absorption band, cm^{-1} , 3) thickness of
absorption layer, mm, 4) concentration, g/ml, 5) width of slit, a) mm,
b) spectral, cm^{-1} , 6) root mean square error of particular concentrations
in the range: %, 7) 1, 2-dichloro ethane, 8) 1, 1, 2-trichloro ethane,
9) 1, 1, 1, 2-tetrachloro ethane, 10) 1, 1, 2, 2-tetrachloro ethane,
11) without solvent

Соединение 1	Температура кипения °C 2	Удельный вес 3	Показатель преломления 4
1,2-дихлорэтан	83,5	$d_{20}^{20} = 1,2563$	$n_D^{20} = 1,4440$
1,1,2-трихлорэтан	113,0	$d_{20}^{20} = 1,4406$	$n_D^{20} = 1,4718$
1,1,1,2-тетрахлорэтан	128,0	$d_{20}^{20} = 1,5539$	$n_D^{20} = 1,4823$
1,1,2,2-тетрахлорэтан	143,5—144,5	$d_{20}^{20} = 1,5955$	$n_D^{20} = 1,4935$

Card. 4/5

Quantitative analysis ...

S/032/61/027/003/010/025
B101/B203

Таблица 2 1. Соединение	2. Полоса поглощения, см ⁻¹	3. Толщина поглощающего слоя, мм	4. Концентрация, г/мл	5. Ширина щели		6. Квадратичная относительная ошибка для определения концентрации в пределах, %	
				2.	3.		
				мм	спектральная см ⁻¹	0.4-5	5-100
7 1,2-дихлорэтан	883	1.35	0.032	0.36	5.3	7.5	1.5
8 1,1,2-трихлорэтан	934	0.45	0.032	0.32	5.7	10.5	1.6
7 1,1,1,2-тетрахлорэтан	960	1.35	0.10	0.30	5.9	7.5	1.5
10 1,1,2,2-тетрахлорэтан	1017	0.22	Без расчёта	0.26	6.2	10.5	2.0

Card 5/5

FLISSKIY, M.M.; VESELOVSKAYA, I.Ye.; DZHAGATSPANYAN, R.V.; CHERNYAVSKAYA, O.V.

Anodic process on graphite in the electrolysis of sodium chloride
in the presence of sulfate ions. Zhur.prikl.khim. 34 no.11:2483-
2487 N '61. (MIRA 15:1)

(Sodium chloride)
(Sulfates)

(Electrolysis)

MARTYNOV, Yu.M.; KORNBLIT, I.I.; SMIRNOVA, N.P.; DZHAGATSPANYAN, R.V.

Determination of metal impurities in silicon tetrachloride
and silicon dioxide by the spectrochemical method.

Zav.lab. 27 no.7:839-842 '61.

(MIRA 14:7)

(Silicon compounds) (Metals--Analysis) (Spectrochemistry)

YAKIMENKO, L.M.; KOKHANOV, G.N.; VESELOVSKAYA, I.Ye.; DZHAGATSPANYAN, R.V.

Investigating the electrochemical behavior of titanium and some
other metals during the electrolysis of chloride solutions. Khim.
prom. no.1:43-47 Ja '62. (MIRA 15:1)
(Titanium—Electric properties) (Chlorides) (Electrolysis)

LEVKOVICH, M.M.; MEZHOV, V.D.; DZHAKELI, T.N.

Production of secondary alkyl sulfates. Khim. i tekhn. topl. 1
masel 6 no. 5:24-29 My '61. (MIRA 14:5)
(Paraffins) (Sulfuric acid)

DZHALALOV, A.D., dotsent; MUMINOV, B.M., ordinator

Dynamics of the gastrocolic reflex in gastric and duodenal ulcer.
Med. zhur. Uzb. no.4:20-23 Ap '61. (MIRA 14:5)

1. Iz kliniki gosital'noy terapii Samarkandskogo gosudarstvennogo
meditsinskogo instituta imeni I.P.Pavlova.
(REFLEXES) (PEPTIC ULCER)

GORYAYEV, M.I.; DZHALILOV, D.R.

Study of the essential oil from needles of the juniper (*Juniperus*
turkestanica kom.) Izv. AN Kazakh. SSR Ser. khim. no. 2:107-113
'60. (MIRA 14:5)

(Essences and essential oils) (Juniper)

15.9205

S/081/62/000/012/058/063
B158/B101

AUTHORS: Dzhagatspanyan, R. V., Zetkin, V. I., Motsarev, G. V.,
Filippov, M. T.

TITLE: Chlorination of silicon-containing monomers and polymers
under the effect of gamma-radiation

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 12, 1962, 612-613
abstract 12P282 (Sb. "Radioakt. izotopy i yadern. izlucheniya
v nar.kh-ve SSSR. V. I.". M., Gostoptekhizdat, 1961,
197-200)

TEXT: Polydimethylsiloxane rubber (I) and polyphenylmethylsiloxane (II)
as well as a number of monomers were chlorinated at 0°C under the action
of γ -radiation (Co^{60} with an activity of 1400 g-equiv of Ra). Chlorination
of I takes place easily and rapidly until the introduction of an average
of two Cl atoms into the chain of the polymer, after which the process
rate falls sharply. In a metal autoclave at both 0°C and 60°C
destruction of the polymer takes place. With chlorination of II (molar
Card 1/2

Chlorination of silicon-containing ... S/081/62/000/012/058/063
B158/B101

ratio of Cl:siloxane = 2:1 and 3:1) substitution and addition chlorination takes place in the aromatic ring. With chlorination of $(CH_3)_3ClSi$ (molar ratio of Cl_2 :silane = 0.51:1) the basic product is a monochlorine derivative; chlorination of 18.6 g of ethyl-trichlorosilane (molar ratio of Cl_2 :silane = 0.35:1) gives 6.5 g of α and β -chloroethyl-trichlorosilanes. Chlorination of methyl-phenyl dichlorosilane results in the formation of $(C_6H_2Cl_3)(OCl_3)SiCl_2$ (b. p. 185-188/10 mm). [Abstracter's note: Complete translation.]

Card 2/2

DZHAGATSPANYAN, R.V.; ZETKIN, V.I.; FEDCHENKO, V.S.

Radiative-chemicalsulfochlorination of polyisobutylene. Plast.-
massy no.5:6-9 '62. (MIRA 15:4)
(Propene) (Chlorosulfonylation) (Radiation)

S/019/62/000/017/003/054
A154/A126

AUTHORS: Dzhagatspanyan, R.V., Yakimenko, L.M., Gershenovich, A.I., Zetkin,
V.I., Pospelov, V.Ye.

TITLE: A method of sulfochlorinating polypropylene

PERIODICAL: Byulleten' izobreteniy, no. 17, 1962, 20

TEXT: Class 12o, 23₀₁. No. 149773 (750759/23-4 of October 30, 1961).
Dependent on Author's Certificate No. 128142. This method of sulfochlorinating
polypropylene is dependent on Author's Certificate No. 128142 and differs in that,
to avoid the use of a solvent at room temperature, the process is carried out in
a fluidized bed of polypropylene.

[Abstracter's note: Complete translation]

Card 1/1

S/844/62/000/000/066/129
D204/D307

AUTHORS: Dzhagatspanyan, R. V., Zetkin, V. I., Motsarev, G. V.
and Filippov, M. T.

TITLE: The chlorination of phenylmethyldichlorosilane (I) and
dimethyldichlorosilane (II) under the action of γ irradi-
ation

SOURCE: Trudy II Vsesoyuznogo soveshchaniya po radiatsionnoy khi-
mii. Ed. by L. S. Polak. Moscow, Izd-vo AN SSSR, 1962,
386-389

TEXT: I was chlorinated in sealed ampoules, at 0 and 20°C, under
 γ irradiation (~120 r/sec, over 15 or 30 min), with molar ratios
(n) of Cl_2 : I equal to 0.25:1, 0.5:1, and 1:1, since polychlori-
nated silanes are of interest in preparing fluorinated Si-contain-
ing monomers and polymers. In contrast to chemically initiated
chlorination of I, the present reaction was one of addition of Cl_2
into the aromatic ring rather than substitution into the methyl

Card 1/2

The chlorination of ...

3/844/62/000/000/066/129
D204/D507

group, the main product being a viscous oil, which by chemical and ir spectroscopic tests proved to be $\text{CH}_3\text{C}_6\text{H}_5\text{Cl}_6\text{SiCl}_2$. A small amount of CH_3 -chlorinated compounds was also formed. No product in which chlorination of CH_3 - and C_6H_5 -groups occurred simultaneously was observed, although it might form in initial mixtures richer in Cl_2 . Silane II was similarly chlorinated at 0°C , with n equal to 0.3:1 and 0.5:1, under 2 min doses of γ rays at 120 r/sec, to give ~30% yields of the monochloride and 5 to ~17% yields of the dichloride, the latter becoming greater with increasing n. There are 4 tables.

ASSOCIATION: NII Goskomiteta, Soveta Ministrov SSSR po khimii
(NII for Chemistry of the State Committee, Council of Ministers of the USSR)

Card 2/2

Radiational sulfo-oxidation ...

S/844/62/000/000/067/123
D204/D307

by total sulfuric and sulfonic acids (H_2SO_4 forms as a by-product) increased linearly with the dose of irradiation, the latter varying from 1 to 4 hours at 110 r/sec. The yields were higher for the nonane (8.7 mg - eqts after 4 hours) than for the undecane (7.04 mg - eqts after 4 hours). There are 3 figures and 2 tables.

ASSOCIATION: NII Goskomiteta, Soveta Ministrov SSSR po khimii (NII for Chemistry of the State Committee, Council of Ministers of the USSR)

Card 2/2

S/844/62/000/000/069/129
D20+/D307

AUTHORS: Yakimenko, L. M., Dzhagatspanyan, R. V., Zetkin, V. I.,
Korolev, B. M. and Maksimov, M. P. (deceased)

TITLE: Chlorine exchange between hexachlorocyclohexane (I) and
carbon tetrachloride, under the action of γ radiation

SOURCE: Trudy II Vsesoyuznogo soveshchaniya po radiatsionnoy khimii.
Ed. by L. S. Polak. Moscow, Izd-vo AN SSSR, 1962, 398-402

TEXT: The behavior was studied of purified α -, β -, γ -, and δ -isomers of I, with CCl_4 labelled with Cl^{36} , under the action of γ -irradiation at 120 r/sec over 1 - 15 hours, at $\sim 30^\circ\text{C}$. The mixtures consisted of 250 mg of I, 8 ml of CCl_4 , and 1 ml of labelled CCl_4 , and the chlorine exchange was assessed by the change in the activity of I (dissolved in 2.5 ml of benzene). No transfer of Cl^{36} took place in the absence of irradiation. The activity of α -I and β -I rose to a constant value of $\sim 1\%$ (arbitrary units) after 5 - 6
Card 1/2

Chlorine exchange between ...

S/844/62/000/000/069/129
D204/D507

hours, whilst that of γ -I increased to a constant 0.6% after 8 - 10 hours. The activity of δ -I was 0.5% after 6 hours. At comparatively low doses of irradiation the processes involved are chiefly the chlorination of I and Cl-exchange, whilst further irradiation allows a 3rd reaction, loss of HCl by I (to form pentachlorocyclohexene), to proceed. The rise of the activity of I might be ascribed to a dynamic equilibrium between these processes, but is more probably due to: (1) radiolysis of CCl_4 to CCl_3 and Cl, formation of C_2Cl_6 and Cl_2 , and recombination to CCl_4 ; (2) $\text{RCl} + \text{Cl} \rightarrow \text{HCl} + \text{R}'\text{Cl}$, $\text{R}'\text{Cl} + \text{Cl} \rightarrow \text{R}''\text{Cl}$, $\text{R}''\text{Cl} + \text{CCl}_3 \rightarrow \text{R}'''\text{Cl}$, $\text{R}'''\text{Cl} + \text{CCl}_4 \rightarrow \text{HCl} + \text{CCl}_3$; (3) $\text{RCl} + \text{Cl} \rightarrow \text{RCl}$ (exchange reaction); (4) $\text{RCl} \rightarrow \text{HCl} + \text{R}'''\text{Cl}$ (dehydrohalogenation). Reaction (4) and chlorination reaction (1) are in agreement with the observed considerable evolution of HCl. There are 2 figures and 1 table.

ASSOCIATION: NII Goskomiteta, Soveta Ministrov SSSR po Khimii (NII for Chemistry of the State Committee, Council of Ministers of the USSR)

Card 2/2

S/076/62/036/008/004/011
B101/B144

AUTHORS: Filippov, M. T., Dzhagatspanyan, R. V., Motsarev, G. V., and
Zetkin, V. I.

TITLE: Infrared spectra of organochlorosilanes containing chlorine
in the organic group

PERIODICAL: Zhurnal fizicheskoy khimii, v. 36, no. 8, 1962, 1751 - 1754

TEXT: IR spectra of $\text{QH}_3\text{C}_6\text{H}_5\text{SiCl}_2$ (I); $\text{CH}_2\text{ClC}_6\text{H}_5\text{SiCl}_2$ (II); $\text{CHCl}_2\text{C}_6\text{H}_5\text{SiCl}_2$ (III); $\text{CCl}_3\text{C}_6\text{H}_5\text{SiCl}_2$ (IV); $(\text{CH}_3)_2\text{SiCl}_2$ (V); $\text{CH}_2\text{ClCH}_2\text{SiCl}_2$ (VI), and $\text{CHCl}_2\text{CH}_2\text{SiCl}_2$ (VII) were studied with the following results: (1) The 3.35 and 3.4 μ bands correspond to the asymmetric and symmetric stretching vibrations of CH in the methyl group. (2) The position of the bands in the range 11-16 μ strongly depends on the degree of chlorination: The 11.76 - 12.7 μ band of V in VI becomes weaker and is shifted toward longer waves; in VII it splits into two bands. (3) The 12.58 μ band of I corresponds to the Si-bound CH_3 group. It changes with the degree of

Card 1/2

Infrared spectra of...

S/076/62/036/006/004/011
B101/B144

chlorination and disappears in IV. (4) The bands of 13-15 μ for I-IV correspond to the C_6H_5 groups. (5) The 15.62 μ band of VI and the 15.38 μ band of II are ascribed to the $SiCH_2Cl$ group. There are no bands in this range for the other compounds. (6) In the case of IV, 11.36 and 11.90 μ bands were observed which appear due to symmetric and asymmetric stretching vibrations of the C-Cl bond in CCl_3 . This was confirmed by the fact that $CCl_3(CH_3)Si(OC_2H_5)_2$ and $(CCl_3)_2Si(OC_2H_5)_2$ also showed bands in the range 11-11.4 μ which were absent in compounds containing no CCl_3 group. There are 5 figures and 2 tables.

SUBMITTED: November 9, 1960

Card 2/2

YAKIMENKO, L.M.; DZHAGATSPANYAN, R.V.; VESELOVSKAYA, I.Ye.; KHODKEVICH, S.D.

Use of platinum-titanium anodes in the chlorine industry.
Khim.prom. no.10:728-735 0 '62. (MIRA 15:12)
(Chlorine industry) (Electrodes, Titanium)
(Electrodes, Platinum)

DZHAGATSPANYAN, Rafael' Vachaganovich; ROMM, Rudol'f Filippovich;
TATOCHENKO, Lev Kirillovich; FINKEL', E.E., red.; KOGAN, V.V.,
tekhn. red.

[Application of radioisotopes to the control of chemical processes]Primenenie radioaktivnykh izotopov dlia kontrolya khimicheskikh protsessov. Moskva, Goskhimizdat, 1963. 343 p.
(MIRA 16:3)

(Radioisotopes—Industrial applications)
(Automatic control)

KOLBASOV, V.I.; BARDENSHTEYN, S.B.; DZHAGATSPANYAN, R.V.; ZAKHAROV, Ye.V.

Quantitative analysis of technical m-chloronitrobenzene by
infrared absorption spectra. Zav.lab. 28 no.11:1326-1327 '62.
(MIRA 15:11)

(Nitrobenzene--Spectra)

PHASE I BOOK EXPLOITATION

SOV/6377

Dzhagatspanyan, Rafael' Vachaganovich, Rudol'f Filippovich Romm, and
Lev Kirillovich Tatochenko

Primeneniye radioaktivnykh izotopov dlya kontrolya khimicheskikh
protssessov (Application of Radioactive Isotopes in the Control of
Chemical Processes) Moscow, Goskhimizdat, 1963. 343 p. Errata
slip inserted. 5000 copies printed.

Ed.: E. E. Finkel'; Tech. Ed.: V. V. Kogan.

PURPOSE: This book is intended for scientific and technical-engineer-
ing personnel working with devices equipped with radioisotopic
sensors.

COVERAGE: The book deals with fundamentals of the theory of devices
equipped with radioisotopic sensors and the use of these devices
for controlling and regulating chemical-engineering processes.
Methods for detection and recording of nuclear radiation and basic
means of protection from such radiation are described. No
Card 1/A2

Application of Radioactive Isotopes (Cont.)

SOV/6377

personalities are mentioned. Lists of references follow each of the three parts of the book.

TABLE OF CONTENTS:

Foreword 6

PART I. PHYSICAL PRINCIPLES OF THE USE OF RADIOACTIVE ISOTOPES

Ch. 1. Origin and Aspects of Nuclear Radiation 9

Ch. 2. Methods for Recording Nuclear Radiation 43

Bibliography to Part I 84

PART II. STANDARD CIRCUIT DIAGRAMS, THEORETICAL
FUNDAMENTALS, AND OPERATING PRINCIPLES OF INDUSTRIAL DEVICES
WITH RADIOISOTOPIC SENSORS

Ch. 3. Principles of the Use of Radiation Detectors in
Radioisotopic Sensors 85

Card 2/42

DZHAGATSPANYAN, R.V.; ZETKIN, V.I.; POSPELOV, V.Ye.; FEDCHENKO, V.S.

Radiation-induced chemical sulfochlorination of polydimethyl-
siloxane. Plast.massy no.2:16-18 '63. (MIRA 16:2)
(Siloxanes) (Chlorosulfonylation) (Radiation)

L 12964-63
 EPR/EMP(j)/EPF(c)/EWT(m)/BDS AFPTC/ASD PS-4/PC-4/Pr-4 RH/WW
 ACCESSION NR: AP3000393 S/0191/63/000/005/0004/0007

AUTHOR: Dzbagatspanyan, R. V.; Zetkin, V. I.; Pospelov, V. Ye.; Fedchenko, V. S. 72

TITLE: Radiochemical sulfochlorination of polystyrene

SOURCE: Plasticheskiye massy⁴, no. 5, 1963, 4-7

TOPIC TAGS: sulfochlorination, polystyrene, chlorine, sulfur dioxide, cobalt sup 60, sulfuryl chloride

ABSTRACT: Improved properties were anticipated from the sulfochlorination of polystyrene, achieved by reacting 1% polystyrene emulsion with chlorine and sulfur dioxide (in molar ratios of 0.22:1 - 4.05:1) dissolved in carbon tetrachloride and subjected to Gamma-radiation from a Co sup 60 source. Over a range of 0 - 55C, the reaction rate increased with increasing temperature to a maximum at 40C. Increasing the total dose of radiation had little effect on the process, which was all but complete within 15-20 minutes. No clear relationship was found between the rate and outcome of the reaction and the molar ratio of the two gases: although the final sulfur content was more dependent than was the chlorine content on the initial ratio, in no case did the final product contain much more than 3% sulfur. Unlike the other polymers, polystyrene could not be sulfochlorinated with sulfuryl chloride. Sulfochlorinated polystyrene had better adhesive qualities (with glass and

Card 1/2

L 12964-63
ACCESSION NR: AP3000593

metals) than polystyrene, a hardness of approximately 0.9 (pendulum apparatus), an impact strength of approximately 50 kg/sec x cm/cm sup 2, and an elasticity in bend of 1 on the NIK scale. Applied without admixture to iron plates, it withstood 6 hours' exposure to 150C. It was, however, less resistant than polystyrene to the action of acids, alkalis, and water. Orig. art. has: 1 figure, 1 formula, 3 tables.

ASSOCIATION: none

SUBMITTED: 00

DATE ACQ: 10Jun63

ENCL: 00

SUB CODE: MA

NO REF SOV: 005

OTHER: 000

Cord 2/2

YAKIMENKO, L.M.; KOKHANOV, G.N.; VESELOVSKAYA, I.Ye.; DZHAGATSPANYAN, R.V.

Investigating the electrochemical behavior of titanium and its alloys
during the electrolysis of chloride solutions. Titan i ego splavy no.
10:168-175 '63. (MIRA 17:1)

KOLBASOV, V.I.; BARDESHTEYN, S.B.; DZHAGATSPANYAN, R.V.

Quantitative analysis of impurities in chloroform from their
infrared absorption spectra. Zav.lab. 29 no.8:938-940 '63.
(MIRA 16:9)

(Chloroform) (Organic compounds--Absorption spectra)
(Chemistry, Analytical-- Quantitative)

VIRIN, L.I.; BUSHINSKAYA, L.M.; DZHAGATSPANYAN, R.V.

Analysis of methallyl chloride by gas-liquid chromatography.
Zav.lab. 29 no.11:1301-1302 '63. (MIRA 16:12)

VESELOVSKAYA, I.Ye.; FLISSKIY, M.M.; DZHAGATSPANYAN, R.V.; MOROCHKO, L.V.

Study of the adsorption of sulfate ion on a graphite anode
under conditions of chloride electrolysis. Zhur. prikl.
khim. 36 no.10:2179-2183 0 '63. (MIRA 17:1)

100-46098

...W. Stokman, R. V.: 1951, . . .

...chlorination of polyethylene under the influence of chlorination

SOURCE: Plasticheskiye massy, no. 10, 1964, 5-8

sulfochlorination, polyethylene gamma radiation, catalytic sulfochlorination, polyethylene sulfide, polyethylene

177. Sulfochlorinated polyethylene has many industrial uses for its properties. Conventionally catalyzed reaction entailed the use of a catalyst. In the present paper, the reaction is initiated by gamma-rays. The effect of the external dose of radiation on the reaction is studied. A CCl_4 solution of a high-pressure polyethylene with a molecular weight of about 20,000 was irradiated with gamma-rays from a ^{60}Co source. The reaction products and their molecular weights were determined. In the experiments, the radiation dose was varied from 0 to 100 Mrads. Sulfochlorinated polyethylene was obtained.

1. The first part of the report is a summary of the work done during the period covered by the report. It is a brief statement of the objectives of the work, the methods used, the results obtained, and the conclusions reached. It is intended to give the reader a general idea of the work and to provide a basis for the more detailed discussion in the following sections.

2. The second part of the report is a detailed description of the work done. It is a full and complete account of the work, including all the details of the methods used, the results obtained, and the conclusions reached. It is intended to provide the reader with a complete understanding of the work and to provide a basis for the more detailed discussion in the following sections.

3. The third part of the report is a discussion of the results obtained. It is a full and complete account of the results, including all the details of the methods used, the results obtained, and the conclusions reached. It is intended to provide the reader with a complete understanding of the results and to provide a basis for the more detailed discussion in the following sections.

4. The fourth part of the report is a conclusion. It is a full and complete account of the conclusions reached, including all the details of the methods used, the results obtained, and the conclusions reached. It is intended to provide the reader with a complete understanding of the conclusions and to provide a basis for the more detailed discussion in the following sections.

N NR: AP4046893

RELATION: none

DATE: 00

ENCL: 00

DATE: 00

NO REP SOV: 005

OTHER: 002

4444-45 EWT(m)/EPF(c)/EWP(j)/EWA(h)/EWA(l) PC-Li/Pr-Li/Pa-Li RM

1. 10/14/20

[illegible]

radiation chlorination of ethyltrichlorosilane, isopropyltrichlorosilane and dimethyldichlorosilane

... 475-476

radiation chlorination, reaction with chlorine, and
chlorination, liquid phase radiation chlorination,
chlorination, methyldichlorosilane, chloromethyl-
trichlorosilane, chloromethyltrichlorosilane, and
chloromethyltrichlorosilane.

The reaction mechanism of radiation-initiated chlorination of poly-chlorosilanes was studied. The rate of radiation-initiated chlorination of ethylchlorosilane by chlorine molar concentration ranging from 0.15 to 0.6 mole/l. was determined. It was found that the chlorination of monochloro derivatives and dichloro derivatives of ethylsilane proceeded at different rates depending on the composition of the system. When reactant ratio was 1:1, the rate of chlorination was higher than when

L 16666-65

NR. AP4044020

...ant ratio = 0.15 or 0.18. Chlorination of methyltrichlorosilane
 at solar ratios ranging from
 and total concentration of
 proportional to the solar ra
 rad sec. had no effect on
 to 3 lower; overall
 polymers derivatives. All the
 mechanism is discussed for the chlorination
 chlorosilane wherein the rate of
 chlorination is determined by the rate of
 and the rate of its initiation
 $\text{CH}_3\text{SiCl}_3 + \text{Cl} \rightarrow \text{CH}_3\text{SiCl}_2\text{Cl} + \text{HCl}$
 about 10 times faster than for
 10-40 increased this chlorination
 activation is about 7500 cal/mole
 the chlorination of dimethyl
 chlorosilane, the rate of the initiation
 is 19 times faster than for
 its energy of activation is
 the ratio in radiation chlorination

NP: AP4044020

initiated chlorination; the same radical
of photochemically and photochemically
radiation chlorination.

Nine

11/11/66, 11/11/66

GC

NR REF SOV: 004

OTHER: 000

ACCESSION NR: AP4034544

S/0020/64/155/005/1163/1166

AUTHORS: Dzhagatspanyan, R.V.; Filippov, M.T.; Motsarev, G.V.; Zetkin, V.I.; Rozenberg, V.R.

TITLE: Radiative chlorination of certain organochlorosilanes and organopolysiloxanes

SOURCE: AN SSSR. Doklady*, .v. 155, no. 5, 1964, 1163-1166

TOPIC TAGS: chlorination, irradiation chlorination, organochlorosilane, organopolysiloxane, chlorination mechanism, polydimethylsiloxane, polyphenylmethylsiloxane, ethyltrichlorosilane, methyltrichlorosilane, dimethyldichlorosilane, phenyltrichlorosilane, phenylmethyldichlorosilane, photochemical chlorination, substitution chlorination, addition chlorination, ionic mechanism, free radical mechanism

ABSTRACT: The mechanisms involved in the chlorination of various organosilane derivatives under the influence of Co^{60} radiation were investigated. A polydimethylsiloxane resin, molecular weight 400,000-500,000, was chlorinated at 0°C as a 4% solution in CCl_4 . After

Card 1/3

ACCESSION NR: AP4034544

chlorination under 4200 rad/min. radiation the chlorine content was 50-55%; optimum reaction time was 15-30 minutes. Total radiation greater than 1.25×10^5 rad did not lead to a higher chlorine content, but promoted degradation of the polymer. By chlorinating polyphenylmethylsiloxane under the same conditions, products containing up to 56.1% chlorine were obtained. About 80% of the chlorine reacted with the aromatic nucleus and 20% replaced hydrogens on a methyl group. Chlorination of ethyltrichlorosilane (molar ratio Cl_2 : $\text{C}_2\text{H}_5\text{SiCl}_2=3:7$) at OC using 900 rad/min gave α - and β -monochloroderivatives in a ratio of 1:1.7, corresponding to results obtained by photochemical chlorination. On chlorinating methyltrichlorosilane and dimethyldichlorosilane the amount of monochloro derivatives in the reaction mixture did not depend on the molar ratio of reagents and the change in the amount of dosage did not influence the products of chlorination. The relative reaction rate of methyltrichlorosilane did not depend on the concentration of chlorine and at OC and 3300 rad/min equaled 0.148 ± 0.030 moles/liter-min. The magnitude is proportional to the square root of the power of dosage. The energy

Card 2/3

ACCESSION NR: AP4034544

of activation is about 7300-6100 cal/mole for the reaction. Phenyltrichlorosilane and phenylmethyldichlorosilane were chlorinated at 0-150C at 5900 and 800 rad/min at 0-20C. The chlorine added to the double bond of the aromatic nucleus giving $C_6H_5Cl_6SiCl_3$ and $C_6H_5Cl_6(CH_3)SiCl_2$. This additive chlorination under radiation is analogous to photochemical chlorination. At 50C, addition chlorination products as well as products of substitution chlorination in the methyl group and the aromatic nucleus were formed. At 100-150C substitution chlorination of the aromatic nucleus predominated indicating ionic mechanism for the arylalkylchlorosilanes. A free radical mechanism was postulated for the alkylchlorosilanes. Orig. art. has: 11 equations and 1 table

ASSOCIATION: None

SUBMITTED: 16Nov63

ENCL: 00

SUB CODE: 00

NR REF SOV: 005

OTHER: 002

Card 3/3

MOTSAREV, G.V.; YAKUBOVICH, A.Ya.; ROZENBERG, V.R.; FILIPPOV, M.T.;
DZHAGATSPANYAN, R.V.; BARDENSHTEYN, S.B.; KOLBASOV, V.I.;
ZETKIN, V.I.

Halogenation of aromatic silanes. Part 17: Addition of chlorine
to phenyl-trichlorosilane. Preparation of hexachlorocyclohexyl-
trichlorosilane and the mechanism of its formation. Zhur. ob.
khim. 35 no.7:1178-1183 J1 '65. (MIRA 18:8)

L 27304-66 EWT(m)/EPF(n)-2/ENP(j)/I/ENA(h)/ENA(1) IJP(z) GG/RM

ACC NR: AP6008980

(A)

SOURCE CODE: UR/0190/65/007/011/1959/1963

AUTHORS: Dzhagatspanyan, R. V.; Kolbasov, V. I.; Bardenshteyn, S. B.; Korolev, B. M.; Romanskiy, I. A.; Zetkin, V. I.

ORG: none

TITLE: The structure of radiation chlorinated and sulfochlorinated polyethylene

SOURCE: Vysokomolekulyarnyye soyedineniya, v. 7, no. 11, 1965, 1959-1963

TOPIC TAGS: polymer, polyethylene, chlorination, aliphatic compound, chlorine

ABSTRACT: The structure of radiation chlorinated and sulfochlorinated polyethylene in the solid state and in solution was studied by IR spectroscopy. The polyethylene specimens were prepared after the method of R. V. Dzhagatspanyan, L. M. Yakimenko, A. I. Gershenovich, and V. I. Zetkin (Avt. svid. No. 150625, 1961; Byull. izobreteniy, 1963, No. 20, 93). The IR spectra of the investigated compounds are presented. It was found that the IR spectra of bulk radiation sulfochlorinated polyethylene were identical to those sulfochlorinated in bulk by chlorine. It is concluded that chlorination of the polymer occurs more readily in the amorphous phase than in the crystalline phase. Orig. art. has: 2 graphs.

SUB CODE: 11/ SUBM DATE: 26Dec64/ ORIG REF: 003/ OTH REF: 005

Card 1/1

UDC: 678.01:53+678.743

DAVIDOVICH, B.V.; YAFIMENKO, I.M.; SINITSYN, V.I.; LYASHIN, Ya.O.;
SINITSYN, V.I.; LUKMAN, B.Ya.

Radiochemical sulfochlorination of kerosene and synthetic. Radiochem.
41 no.4:7-11 Ap '65. (MIRA 18:8)

KHROMENKOV, L.G.; DZHAGATSPANIAN, R.V.; SOKOLOV, V.A.; KOROLEV, B.M.;
ZETKIN, V.I.

Structure formation in radiation sulfochlorinated polyethylene
and its solutions. Vysokom.sped, 7 no.10:1776-1778 0 '65.
(MIRA 18:11)

ZETKIN, V.I.; PANCHENKOV, G.M.; ZAKHAROV, Ye.V.; KOLESNIKOV, I.M.;
DZHAGATSPANYAN, R.V.

Chlorination and sulfochlorination of organic compounds in
apparatus with periodical and continuous action. Khim. prom.
41 no.10:733-734 O '65. (MIRA 18:11)

DZHAGATSPANYAN, R.V.; KOLBASOV, V.I.; BARDENSHTEYN, S.B.; KOROLEV, B.M.;
ROMANSKIY, I.A.; ZETKIN, V.I.

Structure of radiation chlorinated and sulfochlorinated polyethylene.
Vysokom. soed. 7 no.11:1959-1963 N '65. (MIRA 19:1)

1. Submitted December 26, 1964.

VIRIN, L.I.; SAFIN, Yu.A.; TREGER, Yu.A.; DZHAGATSPANIAN, R.V.

Mass spectra and structure of some allyl derivatives. Zhur.fiz.
khim. 39 no.11:2824-2826 N '65.

(MIRA 18:12)

ZETKIN, V.I.; DZHAGATSPANYAN, R.V.; ZAKHAROV, Ye.V.

Chlorination of nitrobenzene. Zhur. prikl. khim. 38 no. 10:
2379-2383 0 '65. (MIRA 18:12)

1. Submitted July 19, 1963.

L 08/09-67 E.M.T(m) WE
ACC NR: AP6015121

(A)

SOURCE CODE: UN/O

0/005/0018/0020

AUTHOR: Dzbagatspanyan, R. V.; Lyashkin, Yu. G.; Filippov, M. Sinitain, V. I.;
Yakimenko, L. M.; Globova, L. I.; Zolkin, V. I.

58

ORG: none

TITLE: Radiation chlorination of kerosene 112

SOURCE: Khimicheskaya promyshlennost', no. 5, 1966, 18-20

TOPIC TAGS: kerosene, gamma radiation, chlorination, photochemistry

ABSTRACT: Groznyy kerosene, from which the aromatic and unsaturated compounds were eliminated by extraction with liquid SO_2 was used during chlorination initiated by γ -radiation of Co^{60} made in the apparatus described by the authors previously (Khim. prom. no. 4, 247, 1965). After purification the kerosene had a molecular weight of 177. Chlorine was passed at the rate of 0.469 g/min in the reactor set into a thermostat with a controlled given temperature. The radiation source was introduced after 15 minutes. The chlorination products were purified from Cl_2 and HCl by passing a flow of nitrogen. The densities and refractive indexes were measured and the degree of chlorination was determined from the graphs, plotted experimentally, showing the dependence of density d_{20}^{20} and the refractory indexes n_D^{20} of the chlorinated products on their chlorine content. Kinetic curves (content of chlorine vs time in min) were

Card 1/2

UDC: 665.634-4 : 66.094.403.085.3

L 08659-67

ACC NR: AP6015121

plotted at various temperatures of chlorination ($T = 20, 40, \text{ and } 60^\circ\text{C}$) and at various doses of radiation ($P = 26.1, 7.3, 1.8, \text{ and } 0.81 \text{ rad/sec}$). The dependence of the radiation-chemical efficiency coefficient G (number of atoms bound with carbon per 100 equivalent) on the radiation dose P was plotted from kinetic curves. The expression $G = 1.22 \cdot 10^3 \cdot \left(\frac{1000}{T} + 5.76 \cdot 10^{-2} [\% \text{Cl}] \right)^{P-0.47}$ well describes the results obtained. (Disagreement of experimental and calculated values averaged $\pm 10.8\%$.) This equation can be used for designing a reactor for a temperature range of $0-100^\circ\text{C}$, a radiation dose of $1-50 \text{ rad/sec}$, and a chlorine content of $5-60\%$. The apparent energy of activation was determined as 3200 cal/mole . The results of radiation chlorination were compared with those of photochemical chlorination and chlorination initiated by azo-bis-isobutyronitril. It was shown that the same degree of chlorination was achieved more rapidly during radiation chlorination. At $T = 20^\circ\text{C}$ and $P = 26 \text{ rad/sec}$, the product containing $\text{Cl} > 60\%$ was obtained in 90 minutes during radiation chlorination. It took 23 and 21 hours to obtain the same product by photochemical chlorination and chlorination initiated by azo-bis-isobutyronitril, respectively. Radiation chlorination also has other advantages: it depends little on temperature and is controlled by the radiation dose (easily controllable rate of chlorination), the rate of the radiation process does not depend on the color of the reacting mixture, and there is a much smaller danger of resinification because of an absence of local overheating. Orig. art. has: 3 fig., 4 formulas, and 1 table.

SUB CODE: 07/ SUBM DATE: none/ ORIG REF: 001/ OTH REF: 001

Card 2/2

L 18417-66 EWT(m)/EWP(f)/T/EWA(h)/EWA(l) RM

ACC NR: AP6003424

SOURCE CODE: UR/0190/66/008/001/0125/0130

AUTHORS: Dzhagatspanyan, R. V.; Bardenshteyn, S. B.; Kolbasov, V. I.; Korolev, B. M.

ORG: none

TITLE: Study of the structure of radiation chlorinated and sulfochlorinated polypropylene 58
19 B

SOURCE: Vysokomolekulyarnyye soyedineniya, v. 8, no. 1, 1966, 125-130

TOPIC TAGS: polymer, polypropylene plastic, polymerization kinetics, IR spectroscopy, spectroscopy, chlorination, organic compound

ABSTRACT: The structure of sulfochlorinated polypropylene, sulfochlorinated by means of radiation in the solid phase, was investigated by IR spectroscopy to extend the work of R. V. Dzhagatspanyan, L. M. Yakimenko, V. I. Zetkin, A. I. Gershenovich, and V. S. Pospelov (Avt. svid. 149773, 1961 g.; RZhKhim, 1963 9T50). A comparison of IR spectra of a specimen chlorinated in solution and in the solid phase is presented. The experimental results are presented graphically (see Fig. 1). It was found that the crystallinity of polypropylene decreases

Card 1/2


UDC: 678.01:53+678.745 2

L 18417-66

ACC NR: AP6003424

0

Fig. 1. Dependence of optical density, measured at the maximum absorption for the band 973 cm^{-1} (measure of isotacticity), on the chlorine content, %.



with increase in the degree of sulfochlorination. The crystallinity of specimens sulfochlorinated in the solid phase is 3.5 times smaller than that of pure polypropylene, and the specimens chlorinated in solution are amorphous. It is concluded that for both types of specimens, i.e., chlorinated in solution and in solid phase, displacement of hydrogen by chlorine takes place more rapidly for CH_2 groups than for CH_3 group hydrogens. Orig. art. has: 8 graphs.

SUB CODE: 11/ SUBM DATE: 26Feb65/ ORIG REF: 003/ OTH REF: 007

Card 2/2 *for*

L 24493-66 EMT(m)/EPF(n)-2/EWP(j)/EWA(h)/EWA(l) IJP(c) GG/RM
 ACC NR: AP6006971 (A) SOURCE CODE: UR/0190/66/008/002/0193/0197

AUTHORS: Dzhagatapanyan, R. V.; Sokolov, V. A.; Khromenkov, L. G.; Korolev, B. M.

ORG: none

TITLE: On x-ray determination of crystallinity in polyethylene, chlorinated and sulfochlorinated by radiation

SOURCE: ¹⁹Vysokomolekulyarnyye soyedineniya, v. 8, no. 2, 1966, 193-197

TOPIC TAGS: polyvinyl chloride, chlorination, polyethylene plastic, x ray analysis, radiation polymerization


ABSTRACT: The degree of crystallinity of polyethylene (I) samples chlorinated and sulfochlorinated in solid phase by penetrating radiation from a Co⁶⁰ source was investigated. The method, involving x-ray study, consists of determining the ratio of the intensities of chlorinated and nonchlorinated samples. This ratio, in turn, gives the ratio of crystalline phases in the samples because only I is in the crystalline phase during the solid phase chlorination. The noninterfering chlorinated products are considered as the amorphous phase. Spectra of chlorinated and nonchlorinated I are shown in Fig. 1. The decrease in peak intensity is a measure

Card 1/3

UDC: 678.01:53+678.743+678.745

L. 24493-66
ACC NR: AP6006971

Fig. 1. Spectra of untreated (1) and chlorinated (2) polyethylene.




of the decrease of crystalline phase and of the increase of x-ray absorption coefficient in the chlorinated sample. Measurements of the densities of chlorinated samples indicate that each sample consists of a mixture of I and of the final

Card 2/3

L 24493-66
ACC NR: AF6006971

chlorination product, polyvinyl chloride.^b The relationship between the density and chlorine content is illustrated in Fig. 2.

Fig. 2. Specific volume of chlorinated samples as a function of chlorine content.



Sulfochlorination in the solid phase has a similar effect. Chlorination of I in a CCl_4 suspension leads to retention of crystallinity to a greater degree. Orig. art. has: 1 table, 4 figures, and 3 equations.

SUB CODE: 07/ SUBM DATE: 04Jan65/ ORIG REF: 003/ OTH REF: 007
Card 3/3 *LC*

TUL'CHINSKIY, V.A.; DZHAGETYANTS, R.M.

For further improvement of the management of the construction
for the transportation industry. Transp.stroi. 9 no.5:9-13
My '59. (MIRA 12:12)

1. Zamestitel' nachal'nika Planovo-proizvodstvennogo upravleniya
(for Tul'chinskiy). 2. Nachal'nik organizatsionno-shtatnogo
otdela Planovo-proizvodstvennogo upravleniya (for Dzhagetyants).
(Construction industry--Management)

TUL'CHINSKIY, V.A.; DZHAGETYANTS, R.M.

Modifying the production capability indices and improving the wage system in building organizations of the Ministry of construction for the Transportation Industry. Transp.stroi. 10 no.4:4-7 Ap '60. (MIRA 13:9)

1. Zamestitel' nachal'nika Planovo-proizvodstvennogo upravleniya Mintransstroya (for Tul'chinskiy). 2. Nachal'nik organizatsionno-shtatnogo otдела Planovo-proizvodstvennogo upravleniya Mintransstroya (for Dzhagetyants). (Wages)

(Construction industry--Management)

BARAYAN, A.A.; DZHAGINOV, A.N.

Effect of the profile of the pipeline route and of small gas
withdrawals on the operating characteristic of a gas pipeline.
Gaz.prom. 6 no.4:37-38 '61. (MIRA 14:3)
(Gas, Natural---Pipelines)